

**(12) PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11)** Application No. **AU 200074284 B2**  
**(10)** Patent No. **782653**

(54) Title  
Composition to be administered through mucous membrane

(51)<sup>6</sup> International Patent Classification(s)  
A61K 009/00

(21) Application No: 200074284 (22) Application Date: 2000.09.13

(87) WIPO No: W001/19335

(30) Priority Data

(31) Number (32) Date (33) Country  
99/11465 1999.09.14 FR

(43) Publication Date : 2001.04.17

(43) Publication Journal Date : 2001.06.14

(44) Accepted Journal Date : 2005.08.18

(71) Applicant(s)  
Capsulis

(72) Inventor(s)  
Sophie Gaubert; Rene Laversanne

(74) Agent/Attorney  
Griffith Hack,GPO Box 4164,SYDNEY NSW 2001

(56) Related Art  
AS IN ISR OF WO 2001/019335, 22/3/01

74234/00

(12) DEMANDE INTERNATIONALE PUBLIÉE EN VERTU DU TRAITÉ DE COOPÉRATION  
EN MATIÈRE DE BREVETS (PCT)(19) Organisation Mondiale de la Propriété  
Intellectuelle  
Bureau international(43) Date de la publication internationale  
22 mars 2001 (22.03.2001)

PCT

(10) Numéro de publication internationale  
WO 01/19335 A2(51) Classification internationale des brevets<sup>1</sup>: A61K 9/00App. 16, 14, Rue des As, F-33600 Pessac (FR). LAVER-  
SASSE, René [FR/FR]; 62, Avenue du Parc d'Espagne,  
F-33600 PESSAC (FR).

(21) Numéro de la demande internationale:

PCT/FR00/02523

(22) Date de dépôt international:

13 septembre 2000 (13.09.2000)

(74) Mandataires: GIRAUD, Françoise etc.; Cabinet Beau de  
Loménie, 158, rue de l'Université, F-75340 Paris Cedex 07  
(FR).

(25) Langue de dépôt:

français

(81) États désignés (national): AU, CA, JP, MX, US.

(26) Langue de publication:

français

(84) États désignés (régional): brevet européen (AT, BE, CH,  
CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT,  
SE).

(30) Données relatives à la priorité:

99/11465 14 septembre 1999 (14.09.1999) FR

Publiée:

Sans rapport de recherche internationale, sera republiée  
dès réception de ce rapport.(71) Déposant (pour tous les États désignés sauf US): CAP-  
SULIS [FR/FR]; 218-228 Avenue du Haut-Lévêque,  
F-33600 Pessac (FR).

(72) Inventeurs; et

(75) Inventeurs/Déposants (pour US seulement): GAUBERT,  
Sophie [FR/FR]; Résidence Les Cottages de la Réserve,En ce qui concerne les codes à deux lettres et autres abrégia-  
tions, se référer aux "Notes explicatives relatives aux codes et  
abréviations" figurant au début de chaque numéro ordinaire de  
la Gazette du PCT.

(54) Title: COMPOSITION TO BE ADMINISTERED THROUGH MUCOUS MEMBRANE

(54) Titre: COMPOSITIONS DESTINÉES A UNE ADMINISTRATION PAR VOIE MUQUEUSE

(57) Abstract: The invention concerns a novel use of onion-structured multilamellar vesicles having a liquid-crystal internal struc-  
ture formed by stacked concentric double layers based on amphiphilic agents with layers of water, aqueous solution or solution of  
polar liquid and wherein is incorporated at least an antigen for making a composition, in particular a pharmaceutical composition,  
and more particularly a vaccine composition, to be administered through a mucous membrane to induce a mucous and/or seral  
systemic response and/or for protecting the system against infection caused by said antigen. The invention also concerns a vaccine  
method through a mucous membrane and a method for producing antibodies, in particular, IgA. The invention is particularly appli-  
cable for intranasal administration.

WO 01/19335 A2

(57) Abrégé: La présente invention concerne une nouvelle utilisation de vésicules multilamellaires à structure en oignon présentant  
une structure interne cristal-liquide formée d'un empilement de bicouches concentriques à base d'agents amphiphiles alternant avec  
des couches d'eau, de solution aqueuse ou de solution d'un liquide polaire et au sein desquelles se trouve incorporé au moins un  
antigène, pour la fabrication d'une composition notamment pharmaceutique et plus particulièrement vaccinale destinée à une admi-  
nistration par voie muqueuse pour induire une réponse muqueuse et/ou systémique sérique et/ou pour protéger l'organisme à l'égard  
de l'infection dont est responsable ledit antigène. L'invention concerne également un procédé de vaccination par voie muqueuse et un  
procédé de production d'anticorps, notamment d'IgA. L'invention s'applique tout particulièrement dans le cas d'une administration  
par voie nasale.

## COMPOSITION TO BE ADMINISTERED THROUGH MUCOUS MEMBRANE

The present invention relates to novel compositions for administration via the mucosa. More particularly, it relates to a pharmaceutical composition, in particular a vaccine composition for administration via the mucosa.

It also relates to a method for producing antibodies, in particular IgA.

A number of definitions are given below:

Administration via the mucosa: non-invasive administration of an antigen to a mucous site, for example:

- naso-pharyngeal area;
- buccal area;
- bronchial tree;
- intestine;
- uro-genital tract;
- inner ear;
- conjunctiva;
- mammary, salivary and lachrymal glands.

Of these different constituents of mucus associated lymphoid tissue (MALT), certain introduction routes are more readily accessible and more readily acceptable: nasal, buccal, gastro-intestinal, rectal and vaginal.

Mucous response: The response of the immune system in the mucosa, characterized by production of IgA (isotype in high concentrations in the mucosa) and IgG and/or a cellular response in the mucous site and lymph nodes.

Systemic response: The response of the generalized immune system resulting from the presence of circulating antibodies (IgG and IgA) and/or a cellular response (T helper or CTL) in the secondary lymphoid organs (spleen, nodes).

Dissemination of response: Installation of a mucous response in compartments other than that used for introduction, because of the circulation and relocation of lymphocytes induced at the

administration site (vaginal IgA after nasal administration, for example).

Adjuvant: The substance added to the antigen to amplify and orientate the specific immune response of that antigen.

Synthetic vector/vectorization: Incorporation of the antigen into or on the surface of a particle or vesicle, to protect the antigen and/or facilitate its capture by competent cells of the immune system and/or to facilitate presentation of the antigen by said cells, and to amplify the immune response. The addition of an empty vector to the free antigen provides little or no amplification.

Until recently, vaccines were produced from dead, attenuated or less virulent micro-organisms. Today's pharmaceutical industry seeks to avoid such an approach, to limit secondary effects, to facilitate production, for safety in administration and for efficacy reasons.

Advances in molecular biology have resulted in the industrial production of sub-units of such micro-organisms, in particular proteins that are termed recombinant proteins, either membrane, nuclear or cytoplasmic proteins. Unfortunately, such sub-units are not in themselves sufficiently immunogenic to supply a sufficient response to vaccination.

They have to be supplemented with adjuvants or vectorized to induce a sufficient response.

Today, the only adjuvants that are acceptable for human medicine are aluminum hydroxide or phosphate, also calcium phosphate. These salts are in the form of a suspension of grains of the aluminum or calcium salt, onto the surface of which the antigen is adsorbed. They have a number of disadvantages: they induce local inflammation reactions and the production of IgE, they are not effective for all antigens and they are incapable of causing CTL type cellular mediation reactions. Thus,

until now they have not been used for administration via the mucosa.

Mucous surfaces are very important, firstly because they are present in all tracts and secondly, because they are the first line of defense against the invasion of pathogenic agents.

Mucous surfaces are protected both by innate or non adaptive defense mechanisms (peristalsis, cilia movements and mucous) and by triggering a cellular immune response and an adaptive humoral response specific to the pathogenic agent, which can be generalized to other lymphoid organs. Mucus associated lymphoid tissue (MALT) is responsible for the specific component.

In this respect, vaccination via the mucosa represents a considerable advantage.

Currently, all vaccines except that for poliomyelitis are injectable, which involves:

- medicalization;
- low acceptance;
- discomfort (fever, injection site pain,...);
- risks of contamination (HIV, hepatitis);
- high costs in developing countries.

Administration via the mucosa would be of advantage for a variety of reasons:

- production costs would be lower due to the less stringent preparation conditions compared with those for an injectable vaccine;
- its non invasive nature, which eliminates problems linked to injection mentioned above.

To this end, it must:

- provide immunity at the mucous administration sites, to strengthen the mucous barrier by dint of a specific response, providing an advantage for infections that can be transmitted via the mucosa (respiratory, genital,...);

- provide systemic immunity and an effective generalized response following easy administration;
- not cause secondary effects (local irritation).

5 A fairly complete review of mucosal immunology can be found in the "Handbook of Mucosal Immunology" edited by Pearay Ogra et al., Academic Press, Inc., San Diego, California, USA.

10 A number of trials and studies have been carried out with a view to developing novel adjuvants and vectors that can induce a mucous and systemic response that is effective as regards protection against the pathogenic agent. Different studies concern the use of live recombinant vectors (attenuated bacteria or viruses),  
 15 synthetic vectors (immunostimulating complexes also known as iscoms, an abbreviation of the expression "immunostimulating complexes", liposomes, microspheres...) or micro-organism toxins (cholera toxin (CT), or heat labile E. Coli toxin, (LT)). However, a large proportion  
 20 of these studies are still in the development stage and the majority of formulations have the following disadvantages:

- high toxicity (ST and LT), and thus are not possible for human application;
- 25 • difficulties in preparation (microspheres);
- low stability (liposomes);
- low efficacy (liposomes).

A number of patents concern administration via the mucosa of antigens that are vectorized or supplemented by  
 30 different systems. The following can be cited:

- European patent EP-A-0 440 289 (Duphar), which describes the use of liposomes mixed with an antigen for vaccination against influenza. The lipids act as an adjuvant, the same result being  
 35 obtained when the antigen is mixed extemporaneously with empty liposomes;

- International patent application WO-A-98/10748 (The School of Pharmacy), using cationic liposomes, in the case of injection or of administration via the mucosa;
- 5 • United States patent US-A-5 679 355 (Proteus), describing non-ionic vesicles, which can be administered parenterally or via the mucosa (especially orally).

Further, a great deal of scientific literature  
 10 describes the use of polymer microparticles as a mucous membrane antigen vector. A good review of that literature can be found in: D. T. O'HAGAN, Adv. Drug. Deliv. Rev., 34 (1198), 305-320.

However, none of the above technologies has yet  
 15 culminated in a commercially available product. Antigen presentation methods must, therefore, be further improved and the development of novel adjuvants or antigen vectors remains an urgent need in the vaccine industry.

Vesicles, spherical objects formed by a molecular  
 20 arrangement of amphiphilic molecules, include multilamellar vesicles with an onion-like structure which have formed the basis of a large amount of research and have given rise to several patents (WO-A-93/19735; WO-A-95/18601; WO-A-97/00623; WO-A-98/02144;  
 25 WO-A-99/16468). They are distinguished from liposomes by the following:

- their preparation method, which starts from a lamellar phase in thermodynamic equilibrium;
- their internal liquid crystal structure formed by  
 30 a regular stack of concentric bilayers of amphiphilic compounds alternating with layers of water or an aqueous solution or a solution of a polar liquid (for example glycerol);
- the varied nature of the amphiphilic molecules  
 35 that can be used to constitute them, used alone or as a mixture.

International patent application WO-A-99/16468 describes vesicles with an onion-like structure incorporating an antigen therein and which can amplify the immune response to that antigen. The examples of that International patent application, which for the first time describes incorporation (also termed encapsulation in that document) of antigens in a multilamellar vesicle with an onion-like structure, clearly show that such encapsulation can substantially amplify the immune response during parenteral administration of an antigen encapsulated in a multilamellar vesicle with an onion-like structure.

The present inventors have now discovered that administration via the mucosa, in particular nasal administration, of compositions of the type described in WO-A-99/16468, causes an immune response not only in different mucous sites but also in the systemic compartment. This response is characterized by immunoglobulin A (IgA) induction. Such a result is not at all obvious having regard to WO-A-99/16468 since, under conditions in which injection was carried out in the examples of that application, even in the encapsulated form, the antigen did not induce any detectable IgA response.

Immunoglobulins (Ig) or antibodies (Ab) associated with cellular components are essential protagonists in the immune response directed against a pathogen. Igs are proteins that are highly specific as regards the antigen.

Different classes of antibodies have been classified, which are distinguished by their mode of induction, their localization and their function (recognition, neutralization of toxic activity or of enzymatic activity).

In the blood compartment, the majority of Ig produced is circulating IgG (sub-divided into different sub-classes), IgA and IgE remaining very much in the minority. In contrast, the major isotype in the mucosa



is IgA, which is produced by IgA plasmocytes of the mucous lymphoid system and secreted actively by the epithelium of the mucosa.

IgA are specialized antibodies and are adapted to defending mucous surfaces that are constantly exposed to pathogens. They are adapted since, because of their biochemical structure (glycosylation, polymerization, secretory piece), they resist the effects of proteases secreted by the micro-organisms and since, in contrast to other isotypes, they limit inflammatory reactions in those compartments in a state of constant activation. They are specialized since they act on a number of levels: secreted, they bind pathogens, limit the effects of toxins and the entry of pathogens by neutralization; they can also act on the epithelium during their transcytosis, or in the lamina propria. Thus, they represent an essential active barrier which, associated with non specific mechanisms, limit invasion by pathogens.

Naturally, cytotoxic and specific responses combine with the antibody response to eliminate the pathogen. If the local response is not sufficient, a systemic response is triggered.

One highly interesting characteristic of the mucous immune system is the re-circulation of B and T lymphocytes induced at a mucous site and their possible domiciliation in sites other than the induction site, propagating the specific response to other tracts. This phenomenon is very important as regards vaccination.

While the mechanisms for induction of a mucous immune response are still in the course of being elucidated, it often remains extremely difficult to induce a mucous response by parenteral administration of an antigen (with or without adjuvant) and even by mucous administration of the antigen. In fact, parenteral administration of an antigen leads to the induction of specific circulating antibodies of the IgG type. This

type of injection cannot induce IgA in the mucous (nor the systemic) compartments. It is very clear that parenteral vaccination will not reinforce the natural local defenses required to combat many respiratory or genital infections, for example.

For these reasons, it is completely surprising even in the light of the teaching of WO-A-99/16468 that the same antigen incorporated in the same vesicles, administered via the mucosa (nasally) can induce not only a systemic response (IgG in serum) confirming the preceding results by another route, but also a response at mucous sites. Incorporation of the antigen into vesicles with an onion-like structure causes the production of IgA in the compartment linked to the administration site (bronchio-pulmonary) but also in other mucous compartments with a predominance in genital secretions. Further, it should be noted that human serum albumin (HSA) which is used by way of example is a very slightly immunogenic antigen which, when administered nasally alone without incorporation into vesicles, is incapable of inducing any response, whether mucous or systemic.

Further, it should be noted that, compared with parenteral administration, administration via the mucosa necessitates loading the antigen into the mucosa and thus penetration of the antigen, which implies that the antigen must be presented in an optimal manner to defeat the non specific defense mechanisms (movement of cilia, mucus) and to resist the enzymes present in the mucosa or on its surface. It appears that these parameters are optimized by the structure of the vesicles during administration via the mucosa. These functions of vesicles in administration via the mucosa are not necessarily present during parenteral administration, where injection directly reaches zones more favorable to capture by cells of the immune system.

The result observed by administration via the mucosa is thus not a simple extrapolation of the result obtained by parenteral administration, but a demonstration of a novel characteristic of vesicles with an onion-like structure.

5 The present inventors have now discovered that vesicles identical to those described in International application WO-A-99/16468 incorporating an antigen cause an immune response that is much stronger than  
10 administering the free antigen. Further, for certain antigens that induce no response by mucous administration, administering them incorporated into multilamellar vesicles with an onion structure can induce a remarkable response. Finally, the induced response is  
15 found not only locally in the mucous compartment (mucous response), but also in the blood circulation (systemic seric response).

Furthermore, these multilamellar vesicles are prepared from biocompatible constituents that are known  
20 to be innocuous. Still further, the preparation method is simple to carry out and requires only routinely used chemical apparatus. The fact that the process uses a lamellar phase initially in thermodynamic equilibrium endows it with excellent reproducibility, and the  
25 vesicles obtained are highly stable.

In a first aspect, the invention provides the use of multilamellar vesicles with an onion-like structure having an internal liquid crystal structure formed by a stack of concentric bilayers based  
30 on amphiphilic agents alternating with layers of water, an aqueous solution or a solution of a polar liquid and into which at least one antigen is incorporated, for the production of a composition, more preferably a pharmaceutical composition such as a vaccine  
35 composition, for administration via the mucosa; wherein the diameter of said vesicles is in the range 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ .

In a second aspect, the invention provides a method of treating a human or animal body by vaccination, characterized in that it comprises administration via the mucosa of a composition as defined in the first aspect of the invention.

In a third aspect, the invention provides a method of producing antibodies, characterized in that it comprises introducing into a host organism, via the mucosa, lamellar vesicles with an onion-like structure having an internal liquid crystal structure formed by a stack of concentric bilayers based on amphiphilic agents alternating with layers of water, an aqueous solution or a solution of a polar liquid and into which at least one antigen is incorporated, wherein the diameter of said vesicles is in the range 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ ; then removing and purifying said antibodies.

In a fourth aspect, the invention provides a method of producing IgA, characterized in that it comprises introducing into a host organism, via the mucosa, lamellar vesicles with a multilamellar onion-like structure with an internal liquid crystal structure formed by a stack of concentric bilayers based on amphiphilic agents alternating with layers of water, an aqueous solution or a solution of a polar liquid and into which the appropriate antigen is incorporated, wherein the diameter of said vesicles is in the range 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ ; and removing and purifying said immunoglobulins.

In a fifth aspect, the invention provides a method of treating a human or animal body by vaccination, characterized in that it comprises administration via the mucosa of a composition comprising:

multilamellar vesicles with an onion-like structure having an internal liquid crystal structure formed by a stack of concentric bilayers based on amphiphilic agents alternating with layers of water, an aqueous solution or a solution of a polar liquid and into which at least one antigen is incorporated; wherein the diameter of said vesicles is in the range 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ .

9b

As used herein, a reference to a "composition of the invention" is a reference to a composition described in the first aspect of the invention. As used herein, a reference to "vesicles of the invention" or "multilamellar vesicles of the invention" is a reference to the vesicles described in the first aspect of the invention.

The term "incorporated", the use of which appears to us to be preferable to that of the term "encapsulated",

10

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210

means that the antigen or antigens form an integral part of the entity constituted by the vesicle. In fact, molecules of antigen(s) can be found in any layer between the center and the periphery of said vesicle.

5 As will become clear from the following detailed description and examples, the pharmaceutical compositions used in the invention allow the preparation of a mucous vaccine intended to induce a mucous response and/or a systemic seric response.

10 As will become clear from the examples, when they are administered via the mucosa, the vesicles with an onion-like structure as defined above incorporating an antigen induce the production of antibodies in man and in animals.

15 As indicated above, one advantage of the invention is to induce a very high production of antibodies characterized by the presence of IgA and IgG. This implies an increase in the frequency of lymphocytes carrying IgA and IgG specific to the antigen. The preparation can thus be used for the purposes of  
20 activation and differentiation of antigen-specific B lymphocytes that can then be used in cell fusion to produce monoclonal antibodies. In fact, specific lymphocytes present in large quantities in nodes draining  
25 the administration site can be immortalized by fusion with a non secreting myeloma and result in hybridomas secreting monoclonal antibodies.

Its capacity to simplify and increase the antibody response means that the invention can also be employed  
30 for the production of antibodies, in particular polyclonal IgA, an isotype that is difficult to generate in conventional operational modes, or polyclonal IgG. These antibodies can be used for non therapeutic purposes, for example for research, more particularly for  
35 biological or immunological research.

Antibody sampling and purification methods are known to the skilled person.

Thus, the invention also concerns a method for producing antibodies, more particularly IgA.

Further, it has been shown that the composition of the invention, when administered via the mucosa, induces  
 5 a protection of the organism to the infection for which the antigen incorporated into said composition is responsible.

As discussed, in a further aspect, the invention concerns a method for treating a human or animal body by  
 10 vaccination via the mucosa, in which a composition is administered that contains multilamellar vesicles with an onion-like structure having an internal liquid crystal structure formed by a stack of concentric bilayers based on amphiphilic agents alternating with layers of water,  
 15 an aqueous solution or a solution of a polar liquid and into which at least one antigen is incorporated; wherein the diameter of the vesicles is in the range 0.1 $\mu$ m to 25  $\mu$ m.

As will become apparent from the following description and examples, it has been shown that when  
 20 administered via the mucosa, the compositions of the invention in which the antigen is incorporated into a vesicle with an onion-like structure as defined above can cause a mucous immune response and stimulate the common mucous lymphoid tissue. Such capacities have been shown  
 25 to be of particular importance regarding antigens with a vaccinating potential as opposed to invasive pathogens with mucous tropism.

The dual possibility of inducing both a mucous response and a systemic response is of huge interest in  
 30 vaccination as it simplifies administration while offering immunity on a number of fronts: mucous to defend against micro-organism access, and systemic to defend against more disseminated or generalized infections.

Further, the results obtained in the context of the  
 35 invention demonstrate that the invention can not only amplify the antibody response but can also generate an immune response with a protective efficacy against

infection. Therefore, the invention is applicable both to the production of antibodies and to vaccination.

The results obtained confirm that vesicles administered via the mucosa, in particular nasally, can be used to induce or amplify the antibody response in the mucous compartments, to disseminate this response to other sites more distant from the administration site, and generate a systemic response.

The diameter of the vesicles used in the invention is in the range 0.1 micrometers ( $\mu\text{m}$ ) to 25  $\mu\text{m}$ , preferably in the range 0.2  $\mu\text{m}$  to 15  $\mu\text{m}$ .

More precisely, the vesicles used in accordance with the invention are preferably constituted by a plurality of layers of amphiphilic agents alternating with layers of aqueous or polar phase. The thickness of each of these layers is molecular, typically of the order of 5 nanometers (nm) to 10 nm. For a stack of ten to a few tens of layers, then, a diameter is obtained that is in the range from 0.1  $\mu\text{m}$  to a few tens of micrometers. This has been observed experimentally, the vesicles being observable under optical microscopes (in polarized light to provide a better contrast because of their birefringence), either as unresolved points for the smallest layers, or as birefringent spheres for the largest. The size profile can be studied using a laser granulometer (using static laser beam diffusion, analyzed under a plurality of angles). In general, a gaussian profile is obtained centered on a value of 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ , showing a slight heterogeneity of size for a given formulation, under the given operating conditions for the preparation.

As described above, vesicles into which the antigen is incorporated have a multilamellar onion-like structure and are constituted, from their center to their periphery, by a succession of lamellar layers separated by a liquid medium. These vesicles can be obtained by a method comprising preparing a lamellar liquid crystal



phase and its transformation by applying shear. Such a method has been described in particular in WO-A-93/19735 claiming priority from French patent FR-A-2 689 418 or WO-A-95/18601, hereby incorporated by reference.

5 In French patent FR-A-2 689 418, this transformation can be carried out during a homogeneous shear step for the liquid crystal phase, which produces vesicles that are also known as microcapsules, with a controlled size. However, adjusting the formulation of the lamellar liquid  
10 crystal phase, in particular the nature of the surfactants forming part of its composition, can enable to transform this liquid crystal phase into vesicles by simple mechanical stress, in particular when mixing the constituents.

15 Such vesicles also have the advantage, inter alia, of being capable of being prepared by a particularly simple preparation method that can employ a wide variety of surfactants.

A further advantage, also linked to the  
20 method used to prepare the vesicles with an onion-like structure used in accordance with the invention, lies in the fact that the active ingredients and additives are incorporated prior to forming the vesicles, which provides an excellent encapsulation yield, and therefore  
25 better efficacy, and also economizes on very expensive molecules.

Such structures are advantageously obtained by incorporating at least one antigen into a lamellar liquid crystal phase comprising at least one surfactant then  
30 transforming this lamellar liquid crystal phase into a dense phase of small multilamellar vesicles.

Thus, the vesicles used in the invention can be obtained using a process in which a liquid crystal lamellar phase incorporating at least one antigen is  
35 prepared and said liquid crystal phase is caused to rearrange itself into multilamellar vesicles by applying shear.

This shear can be homogeneous shear, with the advantage of resulting in perfectly homogeneous vesicles. However, simple mechanical agitation can prove sufficient to result in the formation of the multilamellar vesicles of the invention.

The antigen can be any molecule against which an immune response is desired, whether exogenous such as an infectious pathogenic organism, parasite or micro-organism (yeast, fungus, bacteria or virus), or of intrinsic natural origin (auto-immune disease or cancer). It may have different biochemical natures.

In particular, it may be an antigen selected from the group consisting of :

- proteins, in particular extracted or recombinant proteins, which may or may not be glycosylated;
  - peptides;
  - lipopeptides;
  - polysaccharides;
- or is a mixture of a plurality of these components.

The multilamellar vesicles with an onion-like structure are prepared using methods described above, in particular in International patent application WO-A-99/16468. The amphiphilic molecules used to prepare them will be selected, but this is not obligatory, from molecules that are described in the pharmacopeia, or are already used in drugs applied to the mucosa.

According to an advantageous embodiment, the membranes of the vesicles contained in the compositions of the invention contain at least one surfactant selected from the group consisting of :

- phospholipids, which may or may not be hydrogenated;
- linear or branched, saturated or mono- or poly-unsaturated C<sub>6</sub> to C<sub>30</sub> fatty acids, in the form of

an acid or of a salt of an alkali or alkaline-earth metal, or of an amine;

- esters, which may or may not be ethoxylated, of said fatty acids and
  - 5       • saccharose;
  - sorbitan;
  - mannitol;
  - glycerol or polyglycerol;
  - glycol;
- 10       • mono-, di- or tri-glycerides or mixtures of glycerides of said fatty acids;
- linear or branched, saturated or mono- or poly-unsaturated C<sub>6</sub> to C<sub>30</sub> fatty alcohols, which may or may not be ethoxylated;
- 15       • ethers, which may or may not be ethoxylated, of said fatty alcohols and:
  - saccharose;
  - sorbitan;
  - mannitol;
  - 20       • glycerol or polyglycerol;
  - glycol;
- polyethoxylated vegetable oils, which may or may not be hydrogenated;
- block polymers of polyoxyethylene and polyoxypropylene (poloxamers);
- 25       • polyethyleneglycol hydroxystearate;
- alcohols with a sterol skeleton such as cholesterol or sistosterol;
- sphyngolipids;
- 30       • polyalkylglucosides;
- copolymers of polyethylene glycol and alkylglycol (for example the ELFACOS family from AKZO NOBEL);
- di- or tri-block copolymers of ethers of polyethyleneglycol and polyalkyleneglycol (for
- 35       example the ARLACELL family from ICI).

Optional co-surfactants can be added to these surfactants (which can be used alone or as a mixture), to

improve the rigidity and tightness of the membranes forming the vesicle. Examples of such molecules that can be cited are:

- cholesterol and its derivatives, in particular charged or neutral cholesterol esters such as cholesterol sulfate;
- other derivatives with a sterol skeleton, in particular those of plant origin (sitosterol, stigmasterol,...);
- ceramides.

The formulation can advantageously involve a mixture of surfactant molecules. In general, at least two different surfactants are used with different hydrophilic-lipophilic balances, which allows continuous adjustment of the bilayer properties and can control the appearance of an instability, which governs the formation of the multilamellar vesicles.

Advantageously, from the above surfactants, two surfactants are selected with relatively different properties, in particular a different hydrophilic-lipophilic balance (HLB). The first surfactant will advantageously have a hydrophilic-lipophilic balance in the range 1 to 6, preferably in the range 1 to 4, while the second surfactant will have a hydrophilic-lipophilic balance in the range 3 to 15, preferably in the range 5 to 15.

The preparation obtained after transformation of the lamellar liquid crystal phase into multilamellar vesicles can then be diluted, in particular with an aqueous solvent such as a buffer solution, a saline solution or a physiological solution, for example, to obtain an aqueous suspension of vesicles.

The encapsulation technique used in the present invention can readily achieve very high encapsulation yields, which may be as high as 100%. However, such yields are not always vital to the function of the envisaged applications.

The encapsulation yield of the antigen(s) in the compositions of the invention is advantageously more than 50%, preferably more than 80%.

5 It appears that the structure of the vesicles is responsible for the particularly advantageous results obtained, and that the multilamellar vesicles of the invention allow the antigen to arrive intact at the antigen presenting cells (APC) and assist its capture by these cells. It thus appears that the function of the  
10 vesicles of the invention is to vectorize, protect and improve capture of the antigen by the immune system.

A further advantage of the technology is that natural or artificial polymers can be added to this  
15 formulation, such as polysaccharides (alginates, chitosan, etc.) to reinforce the solidity of the vesicle, and to enable it to remain longer at the administration site or in the organism, thereby delivering the antigen over a longer period. These polymers can either be  
20 incorporated into the vesicle, or be deposited around it in the form of a coating. In this case, the diameter of the vesicle or particle formed from vesicles coated with the polymer matrix is greater than that of the vesicles alone. These polymers can optionally be cross-linked to  
25 further reinforce their solidity.

Furthermore, and this constitutes a further advantage of the technology, the formulation can be completed by adding immuno-modulating molecules (chitosan, interleukines, ...), which have intrinsic  
30 properties that reinforce amplification and orientation of the immune response.

Vesicles incorporating antigens are advantageously prepared in a process consisting of preparing the lamellar phase in a first stage. This is obtained simply  
35 by mixing the ingredients, in an order determined by the scientist according to the miscibilities of each of the constituents. It may be necessary to heat certain pasty

or solid constituents to facilitate their incorporation. In that case, the antigen is preferably added when mixing is complete to avoid subjecting it to too high a temperature. It is also possible to prepare a mixture of all of the constituents except for the antigen or its aqueous solution in the form of a stock mixture, which is used as required to prepare the lamellar phase. The aqueous solution can contain different constituents that ensure its biological compatibility, in particular the buffer mixtures but also the different antigens. The lamellar phase so prepared is then subjected to moderate shear ( $0$  to  $1000\text{ s}^{-1}$ ) for a limited period ( $0$  to  $60$  minutes).

In the majority of cases, shear is obtained directly by the action of the device performing the mixing. For very small quantities, it can be obtained by hand by mixing the preparation using a microspatula in an Eppendorf type tube.

The sheared lamellar phase is then dispersed in a final medium, in general water or a buffer, which may be identical to that used during the preparation of the lamellar phase. This dispersion is advantageously produced at ambient temperature ( $20^{\circ}\text{C}$ - $25^{\circ}\text{C}$ ) by slowly adding the medium to the lamellar phase with constant stirring.

A preservative and optional other additives intended to complete the galenical formulation can be added to the product.

All of the compositions described above comprising at least one antigen incorporated into vesicles with a lamellar onion-like structure have the advantage of being capable of being used for administration via the mucosa and in particular nasally and induce a mucous and/or seric systemic response.

Example I below clearly demonstrates such an effect with HSA.

Figures 1 and 2 accompanying this example summarize the results obtained from different mucous samples (Figure 1) and from animal serum (Figure 2) after nasal administration of the compositions of the invention (group I) compared with those obtained by administration either of compositions in which the same antigen is free (group II) or compositions containing the same vesicles, but empty (group III).

Example II illustrates a process for immunization with FHA. The results obtained in this example are illustrated in Figures 3, 4 and 5, which respectively show:

- Figure 3: antibody responses in different mucous sites;
- Figure 4: antibody responses in serum;
- Figure 5: the pulmonary bacterial load in mice infected with *Bordetella pertussis*.

#### EXAMPLES

##### EXAMPLE 1: PRODUCTION OF HSA-SPECIFIC ANTIBODIES

##### I - Preparation of vesicles containing human serum albumin (HSA)

###### Formulation (percentage by weight)

1. Potassium oleate (FLUKA):	5.0%
2. Lauric alcohol, ethoxylated with 4 ethylene oxides (SEPPIC):	2.0%
3. Lanolin cholesterol (FLUKA):	5.0%
4. Cholesterol 3-sulfate (SIGMA):	2.5%
5. PBS 1x, sterile (LIFE TECHNOLOGIES):	20.0%
6. Phospholipon 90G soya lecithin (NATTERMANN):	45.5%
7. Human albumin (SIGMA), with 30 mg/ml in PBS 1x:	20.0%

###### Operating procedure

The components were sterilized with UV irradiation for 60 minutes. The containers and accessories (spatulas, stirrers...) were flame sterilized immediately prior to use.

Constituents 1 to 5 are introduced into a pill maker in no particular order then heated to 80°C for 60 minutes

with very vigorous magnetic stirring. Total dissolution of constituents 3 and 4 was verified microscopically.

The desired quantities of mixture 1 to 5 were introduced into a sterile 1.5 ml Eppendorf tube at ambient temperature then constituents 6 and 7 were added. The ensemble was homogenized using a sterile needle then left overnight at 4°C.

The preparation was then dispersed to 33.33% in sterile PBS 1x.

## 10 **II- Immunization protocol**

To test the effect of the vesicles of the invention during administration via the mucosa, 6-8 week old BALB/c female mice received, nasally, twice (at D0 and D30), the different preparations described below. Nasal administration necessitated anaesthesia of the animals by a solution, a mixture of Ketamine, Valium and Atropine, injected intraperitoneally. One month after the last immunization, the animals were sacrificed, the serum was collected and samples were taken of the mucous secretions with the exception of the vaginal lavages, which were taken from the live mouse over the three days preceding the end of the experiment.

### Immunization groups

The mice were divided into 4 groups, I to IV, group IV constituting a control group (non immunized mice, also known as naïve mice) and the other groups being subjected to an immunization protocol as defined above using the following products:

- Group I:
- 30 • Encapsulated HSA: 20 µl, per nostril, of the vesicles of the invention incorporating HSA, corresponding to 80 µg of HSA per mouse;
- Group II:
- 35 • HSA: 20 µl, per nostril, of the vesicles of the invention incorporating HSA, corresponding to 80 µg of HSA per mouse;



- Group III:

- Empty vesicles: 20 µl, per nostril, of empty vesicles of the invention.

5        Secretion sampling

All samples were taken with cold solutions (4°C) and were placed in ice as they were taken to limit degradation by proteolytic enzymes.

Bronchio-alveolar lavages:

- 10        The mouse trachea was cannulated using a probe and 750 µl of PBS was injected slowly to prevent a hemorrhage, the lungs were washed 3 times with the same solution, and the sample was then centrifuged to eliminate pulmonary cells and separated into aliquots
- 15        that were stored at -20°C until assay.

Vaginal lavages

- These samples were taken from the non-anaesthetized live mouse by injecting 50 µl of PBS into the vaginal orifice and washing the vagina three times with the same
- 20        solution. This sample was taken over three consecutive days to cover variations in the hormonal cycle of the mouse. The secretions were combined and stored at -20°C.

Intestinal lavages

- The intestine was removed, freed from the mesentera
- 25        and rinsed with water to eliminate external blood. It was then cut longitudinally and incubated in ice in 1 ml of a lavage solution enriched with protease inhibitor. The whole was centrifuged and the supernatant was recovered and frozen at -20°C.

30        Antibody assay

- HSA-specific antibodies present in the serum and secretions were assayed using ELISA to determine the HSA specific IgA and IgG (biotinylated anti-IgA/streptavidine peroxidase, anti-IgG peroxidase). The results are
- 35        expressed as the mean titer determined with respect to a reference serum from the naïve mouse and corresponding to

the inverse of the dilution equal to a reference threshold.

### III - Results

5 The results of the specific antibody assay are shown in the form of two figures (Figure 1 and Figure 2) respectively illustrating the antibody responses obtained in mucous samples for the response associated with the mucosa and in animal serum (systemic response).

10 In these figures, in each case we also indicated the number of mice that reacted to the immunization protocol with respect to the number of mice subjected to this protocol (the mention "n/m" signifying that n mice reacted out of m mice subjected to the immunization protocol).

15 It is clear from these two figures that nasal administration of the vesicles of the invention incorporating HSA cause a substantial production of antibodies in the lungs (Figure 1). Only immunization in the encapsulated form causes a pulmonary immune response of isotype IgA and IgG (the pulmonary existence of these two isotypes has been reported in various publications). While this administration route is not very easy in an animal, all of the animals immunized nasally with the vesicles of the invention responded.

25 Analysis of the other mucous samples revealed the presence of HSA specific IgA in the vagina and intestine of animals immunized by the vesicles of the invention incorporating HSA, which mucosa were very distant from the administration site, with a predilection for vaginal mucosa. These responses indicate a generalization of the response induced in the pulmonary or nasal mucous lymphoid tissue and the circulation and re-distribution of activated and differentiated lymphocytes close to the administration site.

35 In contrast to the pulmonary mucosa, the predominant HSA specific isotype in the intestine and vagina was IgA. While IgG<sub>H+L</sub> titers could not be carried out for the

vaginal lavages, similar experiments carried out with another antigen indicated the predominance of IgA in vaginal secretions.

Only the vectorized form of the antigen resulted in an intense and specific response. The few animals presenting a detectable IgA response in the groups immunized with free HSA (titer of 3 as opposed to 650 in the vaginal secretions with vesicles of the invention containing HSA) could not be differentiated from the group using empty vesicles of the invention.

Studying the animal serum (Figure 2) indicates that the vesicles of the invention incorporating nasally administered HSA are capable of causing a systemic response characterized by the major presence of IgG but also a large quantity of seric IgA. Only the vectorized form led to this production (IgA, HSA alone = 35; IgA, vesicles of the invention HSA = 17620). It is remarkable to obtain a systemic response to the vectorized antigen. It should be remembered that it is extremely difficult to induce a seric IgA response by systemic immunization with an antigen alone or accompanied by an adjuvant authorized for human use.

In conclusion, not only does the encapsulated form cause a pulmonary immune response but also, only this form permits dissemination of the induced response in the respiratory tract to other mucosa, indicating that the vesicles of the invention are powerful vectors for inducing mucous immunity, encouraging loading and induction of the response at a site and re-distributing it to other sites.

#### **EXAMPLE II: PRODUCTION OF FHA-SPECIFIC ANTIBODIES AND PROTECTION AGAINST *BORDETELLA PERTUSSIS***

FHA protein is a filamentous adhesin of *Bordetella pertussis*, the whooping-cough bacterium. In contrast to the HSA used above, FHA is more immunogenic and is thus susceptible of itself inducing an antibody response. This protein forms part of protective antigens contained

in commercial vaccines against whooping-cough. Finally, the murine model can be infected with *Bordetella pertussis* and it is thereby possible to carry out an infection test following mucous administration and to demonstrate the protective nature of the response induced by immunization.

#### A - Antibody induction

##### I - Preparation of vesicles containing FHA

###### Formulation

1. Potassium oleate (FLUKA):	5.0%
2. Lauryl ether of PEG-4 (Simulsol P4, SEPPIC):	2.0%
3. Lanolin cholesterol (FLUKA):	5.0%
4. Cholesterol 3-sulfate (SIGMA):	2.5%
5. PBS 1x, sterile (LIFE TECHNOLOGIES):	20.0%
6. Soya lecithin (Phospholipon 90G NATTERMANN):	45.5%
7. FHA from <i>Bordetella pertussis</i> in PBS (1 mg/ml):	20.0%

The operating protocol for the preparation was similar to that of Example 1. The FHA concentration of the finished product was 3 µg for 40 µl.

##### II - Immunization protocol

To study the antibody response, the mucous immunization and sampling protocol was identical to that described in Example 1. Two groups of 5 animals were constituted, group I receiving the antigen encapsulated in the vesicles, and group II receiving the non encapsulated antigen in solution in PBS. For each immunization, each animal received 3 µg of FHA, distributed into two instillations, one into each nostril.

###### Secretion sampling

The samples were taken in a manner that was strictly identical to that described for Example 1.

###### Antibody assay

The FHA specific antibodies present in the serum and secretions were assayed using ELISA using a protocol identical to that described in Example 1, with reagents specific to the FHA antigen.

### III - Results

The results are presented in Figures 3 and 4, showing the mean titers of IgA and IgG antibodies in mucous secretions (Figure 3) and in the serum (Figure 4) for the two groups I and II.

The mucous secretions were respectively studied as described in Example 1, from bronchio-alveolar lavages (A), intestinal lavages (B) and vaginal lavages (C).

In group I, mice immunized by the antigen incorporated into the vesicles of the invention (Figure 3), all of the animals responded to the antigen while when the free antigen was administered, only 1/5 of the animals responded, and then very weakly.

As was the case with HSA in Example 1, a very strong amplification of the antibody response in the mucous sites was observed. Nasal immunization with the antigen incorporated into the vesicles of the invention enables to induce a mucous response (IgA) in the lungs, close to the administration site (IgA titer for an antigen incorporated into the vesicles of the invention, 8925, compared with 4.25 for the free antigen), but also a mucous response disseminated into other secretory compartments (intestine and vagina) (titer of vaginal IgA, 13500 for the antigen incorporated into the vesicles of the invention compared with 2.5 for the free antigen). Further, IgG were detected in significant amounts in the three mucous compartments being studied.

As was the case with HSA, we observed large titers of IgA and IgG antibodies in the circulation (Figure 4) following nasal administration of the antigen incorporated into the vesicles (IgA titer for the antigen in the vesicles of the invention, 1300 compared with 39.2 for the free antigen; IgG titer 1 265 000 compared with 3 000).

The results of this Example 2 completely confirm those obtained for HSA. They confirm that administered nasally vesicles can be used to induce or amplify the

antibody response in mucous compartments, to disseminate this response to other sites more distant from the administration site and to generate a systemic response.

#### **B - Protection test**

5 To check whether the immunization obtained was protective, a protection test known as a challenge test was carried out. In this experiment, immunized mice were infected nasally with a fixed amount of *Bordetella pertussis*, then the animals were sacrificed at different  
10 times, to observe the progress of infection. The measurement was made by counting the colonies in the lungs.

#### **I - Preparation of vesicles containing FHA**

15 The formulation for and preparation of the vesicles were strictly identical to that used to characterize the antibody response.

#### **II - Immunization protocol**

The immunization protocol and the doses of  
20 administered antigens were identical to those used to induce antibody.

Three groups of mice were used:

- I. mice immunized by the antigen incorporated into the vesicles of the invention;
- 25 II. mice immunized by the free antigen in solution in PBS;
- III. non-immunized mice.

Each group was composed of 4 mice, immunized at the same time, with the same batch of antigen, in order to  
30 carry out the same infection measurements under the same conditions.

#### **III - Infection protocol**

Four weeks after the second immunization, the mice were anaesthetized and infected with *Bordetella pertussis*  
35 nasally ( $5 \times 10^7$  bacteria/mouse, administered in 20  $\mu$ l) and the initial infectious load was verified from 3 hours after infection.

#### IV - Protocol for measuring the infectious charge

The mice were sacrificed and their lungs were removed at different times after infection and dilacerated in PBS to obtain a homogeneous suspension.

- 5 Different dilutions of the suspension were spread on specific nutrient medium for *Bordetella* and the colonies were counted after growing for 3 days.

#### V - Results

- 10 The results are shown in Figure 5, which indicates the number of colonies of *Bordetella pertussis* in the lungs of mice 3 days (D3, solid bar) and 5 days (D5, open bar) after infection, in millions of bacteria colonies for the three groups of mice shown below:

- 15 Group I: mice immunized with antigen incorporated into the vesicles of the invention.

Group II: mice immunized with free antigen in solution in PBS.

Group III: non-immunized mice.


- 20 It can be seen that, from the third day following infection, the bacterial load for mice immunized with the antigen incorporated into the vesicles of the invention was lower (factor 2) than that for non immunized mice (factor 8). Remarkably, while 5 days after infection, the mice immunized with the free antigen and the non-  
25 immunized mice experienced the bacterial load increase, the mice immunized with the antigen incorporated into the vesicles of the invention had fewer bacteria on D5 than on D3, indicating favorable evolution of the disease. Further, at this stage, the number of bacterial colonies  
30 in the group immunized with the antigen incorporated into the vesicles of the invention was 6 times lower than the group immunized with free FHA and 13 times lower than in the non immunized group, indicating good protection.

- 35 These results demonstrate that the invention can not only amplify the antibody response, but that it also generates an immune response with protective efficacy

against infection. The invention is thus also applicable to the production of antibodies and to vaccination.

- 5           It is to be understood that a reference herein to a prior art document does not constitute an admission that the document forms part of the common general knowledge in the art in Australia.

- 10           In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated  
15 features but not to preclude the presence or addition of further features in various embodiments of the invention.





THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Use of multilamellar vesicles with an onion-like structure having an internal liquid crystal structure formed by a stack of concentric bilayers based on  
5 amphiphilic agents alternating with layers of water, an aqueous solution or a solution of a polar liquid and into which at least one antigen is incorporated, for the production of a composition for administration via the mucosa; wherein the diameter of said vesicles is in the  
10 range 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ .
2. Use according to claim 1, characterized in that said composition is a composition intended to induce a mucous response.  
15
3. Use according to claim 1 or claim 2, characterized in that said composition is a composition intended to induce a systemic seric response.
- 20 4. Use according to any one of claims 1 to 3, characterized in that said composition is a composition intended to induce the production of antibodies.
5. Use according to any one of claims 1 to 4,  
25 characterized in that said composition is a pharmaceutical composition intended to induce protection of the organism against the infection for which said antigen is responsible.
- 30 6. Use according to claim 5, characterized in that said pharmaceutical composition is a vaccine.
7. Use according to any one of claims 1 to 6,  
35 characterized in that said antigen is of exogenous or intrinsic natural origin.

8. Use according to any one of claims 1 to 7,  
characterized in that said antigen is selected from the  
group consisting of :

- proteins, which may or may not be glycosylated;
- 5   • peptides;
- lipopeptides;
- polysaccharides;
- or is a mixture of a plurality of these components.

10   9. Use according to claim 8, characterized in that said  
antigen is an extracted or recombinant protein, which  
may or may not be glycosylated.

15   10. Use according to any one of claims 1 to 9,  
characterized in that said vesicles contain at least one  
surfactant selected from the group consisting of :

- phospholipids, which may or may not be  
hydrogenated;
- linear or branched, saturated or mono- or poly-  
20   unsaturated C<sub>6</sub> to C<sub>30</sub> fatty acids, in the form of  
an acid or of a salt of an alkali or alkaline-  
earth metal, or of an amine;
- esters, which may or may not be ethoxylated, of  
said fatty acids and
  - 25   • saccharose;
  - sorbitan;
  - mannitol;
  - glycerol or polyglycerol;
  - glycol;
- 30   • mono-, di- or triglycerides or mixtures of  
glycerides of said fatty acids;
- linear or branched, saturated or mono- or poly-  
unsaturated C<sub>6</sub> to C<sub>30</sub> fatty alcohols, which may or  
may not be ethoxylated;
- 35   • ethers, which may or may not be ethoxylated, of  
said fatty alcohols and:
  - saccharose;
  - sorbitan;

- mannitol;
- glycerol or polyglycerol;
- glycol;
- polyethoxylated vegetable oils, which may or may not be hydrogenated;
- block polymers of polyoxyethylene and polyoxypropylene (poloxamers);
- polyethyleneglycol hydroxystearate;
- alcohols with a sterol skeleton;
- sphingolipids;
- polyalkylglucosides;
- copolymers of polyethylene glycol and alkylglycol;
- di- or tri-block copolymers of ethers of polyethyleneglycol and polyalkyleneglycol.

11. Use according to claim 10, characterized in that said vesicles contain cholesterol or sistosterol as the surfactant.

12. Use according to any one of claims 1 to 11, characterized in that said vesicles also contain at least one co-surfactant intended to improve the rigidity and/or tightness of the membranes of said vesicles.

13. Use according to claim 12, characterized in that said co-surfactant is selected from the group consisting of formed by:

- cholesterol and its derivatives;
- derivatives with a sterol skeleton;
- ceramides.

14. Use according to claim 13, characterized in that said co-surfactant is a charged or neutral cholesterol ester.

15. Use according to claim 14, characterized in that said co-surfactant is cholesterol sulfate.

16. Use according to claim 13, characterized in that said co-surfactant is a derivative with a sterol skeleton of plant origin.

5

17. Use according to claim 16, characterized in that said derivative with a sterol skeleton of plant origin is sitosterol or stigmasterol.

10 18. Use according to any one of claims 1 to 17, characterized in that said vesicles also contain an immuno-modulating substance.

15 19. Use according to any one of claims 1 to 18, characterized in that the diameter of said vesicles is in the range 0.2  $\mu\text{m}$  to 15  $\mu\text{m}$ .

20 20. Use according to any one of claims 1 to 19, characterized in that the bilayers of said vesicles comprise at least two surfactants, one of which has a hydrophilic-lipophilic balance (HLB) in the range 1 to 6, and the other has a hydrophilic-lipophilic balance (HLB) in the range 3 to 15.

25 21. Use according to claim 20, characterized in that one of the said surfactants has a hydrophilic-lipophilic balance (HLB) in the range 1 to 4.

30 22. Use according to claim 20 or 21, characterized in that one of the said surfactants has a hydrophilic-lipophilic balance (HLB) in the range 5 to 15.

35 23. Use according to any one of claims 1 to 22, characterized in that the encapsulation yield of the antigen (or antigens) in said vesicles is more than 50%.

24. Use according to claim 23, characterized in that said encapsulation yield is more than 80%.

25. Use according to any one of claims 1 to 24, characterized in that said administration via the mucosa is nasal administration.

5

26. A method of treating a human or animal body by vaccination, characterized in that it comprises administration via the mucosa of a composition as defined in any one of claims 1 to 24.

10

27. A method according to claim 26, characterized in that said administration is carried out nasally.

28. A method of producing antibodies, characterized in that it comprises introducing into a host organism, via the mucosa, lamellar vesicles with an onion-like structure having an internal liquid crystal structure formed by a stack of concentric bilayers based on amphiphilic agents alternating with layers of water, an aqueous solution or a solution of a polar liquid and into which at least one antigen is incorporated, wherein the diameter of said vesicles is in the range 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ ; then removing and purifying said antibodies.

29. A method of producing IgA, characterized in that it comprises introducing into a host organism, via the mucosa, lamellar vesicles with a multilamellar onion-like structure with an internal liquid crystal structure formed by a stack of concentric bilayers based on amphiphilic agents alternating with layers of water, an aqueous solution or a solution of a polar liquid and into which the appropriate antigen is incorporated, wherein the diameter of said vesicles is in the range 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ ; and removing and purifying said immunoglobulins.

30. A method of treating a human or animal body by vaccination, characterized in that it comprises administration via the mucosa of a composition comprising:

5 multilamellar vesicles with an onion-like structure having an internal liquid crystal structure formed by a stack of concentric bilayers based on amphiphilic agents alternating with layers of water, an aqueous solution or a solution of a polar liquid and into which at least one  
10 antigen is incorporated; wherein the diameter of said vesicles is in the range 0.1  $\mu\text{m}$  to 25  $\mu\text{m}$ .

31. Method according to claim 30, characterized in that said composition is a composition intended to induce a  
15 mucous response.

32. Method according to claim 30 or claim 31, characterized in that said composition is a composition intended to induce a systemic seric response.  
20

33. Method according to any one of claims 30 to 32, characterized in that said composition is a composition intended to induce the production of antibodies.

25 34. Method according to any one of claims 30 to 33, characterized in that said composition is a pharmaceutical composition intended to induce protection of the organism against the infection for which said antigen is responsible.

30 35. Method according to claim 34, characterized in that said pharmaceutical composition is a vaccine.

36. Method according to any one of claims 30 to 35, characterized in that said antigen is of exogenous or  
35 intrinsic natural origin.

37. Method according to any one of claims 30 to 36,  
characterized in that said antigen is selected from the  
group consisting of :

- proteins, which may or may not be glycosylated;
  - peptides;
  - lipopeptides;
  - polysaccharides;
- or is a mixture of a plurality of these components.

38. Method according to claim 37, characterized in that  
said antigen is an extracted or recombinant protein,  
which may or may not be glycosylated.

39. Method according to any one of claims 30 to 38,  
characterized in that said vesicles contain at least one  
surfactant selected from the group consisting of :

- phospholipids, which may or may not be hydrogenated;
- linear or branched, saturated or mono- or poly-unsaturated C<sub>6</sub> to C<sub>30</sub> fatty acids, in the form of an acid or of a salt of an alkali or alkaline-earth metal, or of an amine;
- esters, which may or may not be ethoxylated, of said fatty acids and
  - saccharose;
  - sorbitan;
  - mannitol;
  - glycerol or polyglycerol;
  - glycol;
- mono-, di- or triglycerides or mixtures of glycerides of said fatty acids;
- linear or branched, saturated or mono- or poly-unsaturated C<sub>6</sub> to C<sub>30</sub> fatty alcohols, which may or may not be ethoxylated;
- ethers, which may or may not be ethoxylated, of said fatty alcohols and:
  - saccharose;
  - sorbitan;

- mannitol;
- glycerol or polyglycerol;
- glycol;
- polyethoxylated vegetable oils, which may or may not be hydrogenated;
- block polymers of polyoxyethylene and polyoxypropylene (poloxamers);
- polyethyleneglycol hydroxystearate;
- alcohols with a sterol skeleton;
- sphingolipids;
- polyalkylglucosides;
- copolymers of polyethylene glycol and alkylglycol;
- di- or tri-block copolymers of ethers of polyethyleneglycol and polyalkyleneglycol.

40. Method according to claim 39, characterized in that said vesicles contain cholesterol or sistosterol as the surfactant.

41. Method according to any one of claims 30 to 40, characterized in that said vesicles also contain at least one co-surfactant intended to improve the rigidity and/or tightness of the membranes of said vesicles.

42. Method according to claim 41, characterized in that said co-surfactant is selected from the group consisting of formed by:

- cholesterol and its derivatives;
- derivatives with a sterol skeleton;
- ceramides.

43. Method according to claim 42, characterized in that said co-surfactant is a charged or neutral cholesterol ester.

44. Method according to claim 43, characterized in that said co-surfactant is cholesterol sulfate.



45. Method according to claim 42, characterized in that said co-surfactant is a derivative with a sterol skeleton of plant origin.

5

46. Method according to claim 45, characterized in that said derivative with a sterol skeleton of plant origin is sitosterol or stigmasterol.

10 47. Method according to any one of claims 30 to 46, characterized in that said vesicles also contain an immuno-modulating substance.

15 48. Method according to any one of claims 30 to 47, characterized in that the diameter of said vesicles is in the range 0.2  $\mu\text{m}$  to 15  $\mu\text{m}$ .

20 49. Method according to any one of claims 30 to 48, characterized in that the bilayers of said vesicles comprise at least two surfactants, one of which has a hydrophilic-lipophilic balance (HLB) in the range 1 to 6, and the other has a hydrophilic-lipophilic balance (HLB) in the range 3 to 15.

25 50. Method according to claim 49, characterized in that one of the said surfactants has a hydrophilic-lipophilic balance (HLB) in the range 1 to 4.

30 51. Method according to claim 49 or 50, characterized in that one of the said surfactants has a hydrophilic-lipophilic balance (HLB) in the range 5 to 15.

35 52. Method according to any one of claims 30 to 51, characterized in that the encapsulation yield of the antigen (or antigens) in said vesicles is more than 50%.

53. Method according to claim 52, characterized in that said encapsulation yield is more than 80%.

54. Method according to any one of claims 30 to 53,  
characterized in that said administration via the mucosa  
is nasal administration.

5


55. Use according to claim 1 substantially as herein  
described with reference to any one of the Examples.

56. Method according to any one of claims 26, 28, 29  
10 and 30 substantially as herein described with reference  
to any one of the Examples.

Dated this 14th day of June 2005

15 CAPSULIS

By its Patent Attorneys  
GRIFFITH HACK






FIG.1




FIG.2

1/3




FIG.3

FIG.4




FIG.5

